

# LOOK AT HOW WE'VE CHANGED



For less  
cramping,  
fullness, and  
overall discomfort.

Still only 2 LITERS.  
Now with only 2 TABLETS.

## HalfLytey<sup>®</sup> & Bisacodyl tablets Bowel Prep Kit

PEG-3350, sodium chloride, sodium bicarbonate  
and potassium chloride for oral solution and  
bisacodyl delayed-release tablets

HalfLytey and Bisacodyl Tablets Bowel Prep Kit is a gastrointestinal lavage indicated for cleansing of the colon as a preparation for colonoscopy in adults. Most common adverse reactions (<3%) are abdominal pain/cramping, nausea, vomiting and headache. Use is contraindicated in the following conditions: known allergies to polyethylene glycol or other components of the kit, gastrointestinal (GI) obstruction, bowel perforation, toxic colitis, toxic megacolon. Use with caution in patients using concomitant medications (such as diuretics) that increase the risk of electrolyte abnormality, patients with known or suspected hyponatremia, patients with severe ulcerative colitis, ileus or gastric retention. There have been reports of ischemic colitis in patients with use of HalfLytey and 20 mg Bisacodyl Tablets Bowel Prep Kit. However, a causal relationship has not been established. If patients develop severe abdominal pain or rectal bleeding, patients should be evaluated as soon as possible. Patients with impaired water handling who experience severe vomiting should be closely monitored including measurement of electrolytes. Hives and skin rashes have been reported with PEG-based products which are suggestive of an allergic reaction.

*Please see brief summary of prescribing information on adjacent page.*

# HalfLyte<sup>®</sup> & Bisacodyl tablets Bowel Prep Kit

PEG-3350, sodium chloride, sodium bicarbonate  
and potassium chloride for oral solution and  
bisacodyl delayed-release tablets

## REFORMULATION

**results in less cramping, fullness, and overall discomfort.**

- Dosage of bisacodyl is reduced from 20 mg (4 tablets) to 10 mg (2 tablets)

**Still only 2 LITERS.**

**Now with only 2 TABLETS.**

Percentage of Patients Reporting in their Diaries “bothersome” to “severely distressing” Symptoms in Controlled Clinical HalfLyte and Bisacodyl Tablets Bowel Prep Kit Trial\*\*†

	HalfLyte and 10 mg Bisacodyl Tablets Bowel Prep Kit (N=222)	HalfLyte and 20 mg Bisacodyl Tablets Bowel Prep Kit (N=223)
Nausea	13%	21%
Abdominal cramping	7%	14%
Abdominal fullness	11%	13%
Vomiting	5%	8%
Overall Discomfort	14%	20%

\* Patients were specifically asked about the occurrence of the following symptoms: nausea, abdominal cramping, fullness, vomiting, and overall discomfort [see Table 2, full prescribing information].

† The HalfLyte and 20 mg Bisacodyl Tablets Bowel Prep Kit does not provide any additional efficacy benefits over the HalfLyte and 10 mg Bisacodyl Tablets Bowel Prep Kit.

**Brief Summary:** Before prescribing, please see full prescribing information for HalfLyte and Bisacodyl Tablets Bowel Prep Kit. **INDICATIONS AND USAGE:** A gastrointestinal lavage indicated for cleansing of the colon as a preparation for colonoscopy in adults. **CONTRAINDICATIONS:** Patients with the following conditions: known allergies to polyethylene glycol or other components of the kit, gastrointestinal (GI) obstruction, bowel perforation, toxic colitis, toxic megacolon. **WARNINGS AND PRECAUTIONS:** Use with caution in patients using concomitant medications (such as diuretics) that increase the risk of electrolyte abnormality, patients with known or suspected hyponatremia, patients with severe ulcerative colitis, ileus or gastric retention. There have been reports of ischemic colitis in patients with use of HalfLyte and 20 mg Bisacodyl Tablets Bowel Prep Kit. However, a causal relationship has not been established. If patients develop severe abdominal pain or rectal bleeding, patients should be evaluated as soon as possible. Patients with impaired water handling who experience severe vomiting should be closely monitored including measurement of electrolytes. Hives and skin rashes have been reported with PEG-based products which are suggestive of an allergic reaction. **Pediatric Use:** Safety and effectiveness in pediatric patients has not been established. **ADVERSE REACTIONS:** Most common adverse reactions (<3%) are abdominal pain/cramping, nausea, vomiting and headache. **Oral Administration:** Take two bisacodyl tablets with water (do **NOT** chew or crush). Wait for a bowel movement (or maximum of 6 hours) then drink the 2 liter solution at a rate of 8 ounces every 10 minutes. Drink all of the solution. **STORAGE:** Store at 20-25°C (59-86°F). When reconstituted you may keep the solution refrigerated. Discard within 48 hours. **Rx only.** Distributed by Braintree Laboratories, Inc. Braintree, MA 02185.

For additional information, please call 1-800-874-6756 or visit us at [www.halflyte.com](http://www.halflyte.com)