

PRE-PEN[®]

(benzylpenicilloyl polylysine injection USP)

NDC: 49471-001-05

SKIN TEST ANTIGEN

It is estimated that 10% of the US population (30 Million patients) self-report as being penicillin allergic; however, only 10% (3 Million patients) have a true penicillin allergy.¹

Determining the true presence of penicillin allergy can be a critical element in patient care. PRE-PEN is the only FDA approved skin test for the diagnosis of penicillin allergy and is now available from ALK.

Discover the **VALUE** of penicillin allergy testing

- Combat drug resistance
- Provide more cost effective treatments for patients
- Provide access to an additional diagnostic tool



Visit www.pre-pen.com to learn more and view the package insert.

Contact **ALK** to place your **PRE-PEN** order today.



www.alk-abello.com/us



800-325-7354



customerservices@alk-abello.com



888-329-2551

Risk Information:

The risk of sensitization to repeated skin testing with PRE-PEN is not established. Rarely, a systemic allergic reaction including anaphylaxis may follow a skin test with PRE-PEN. To decrease the risk of a systemic allergic reaction, puncture skin testing should be performed first. Intradermal skin testing should be performed only if the puncture test is entirely negative. PRE-PEN is contraindicated in those patients who have exhibited either a systemic or marked local reaction to its previous administration. Patients known to be extremely hypersensitive to penicillin should not be skin tested. Please see the package insert for full prescribing information at www.pre-pen.com.

¹Simpson M.D., Alyson B., et al. "e Acceptability of a Four-Part Protocol for Penicillin Allergy Testing by Practicing Allergists."